

**RECOGNISING PMDD
IN PATIENTS:**

**THE
PMDD
PROJECT**

A GUIDE FOR GPs

Premenstrual Dysphoric Disorder (PMDD) is often misunderstood. As a GP, you play a vital role in ensuring that patients receive an accurate diagnosis.

At present, individuals experiencing PMDD have to wait around 12 years on average to receive a diagnosis. PMDD affects an estimated 1 in 20 menstruating individuals and can be confused with other conditions, such as bipolar disorder, depression, or even severe PMS.

Each patient's experience with PMDD is unique, and their symptoms can vary in intensity and presentation. Additionally, anxiety that often accompanies PMDD may prevent patients from fully expressing the severity of their symptoms, so it's crucial to approach the diagnosis with empathy and understanding.

UNDERSTANDING PMDD

Key characteristics:

PMDD is more than just "bad PMS".

While both involve physical and emotional symptoms before menstruation, PMDD's symptoms are far more severe, particularly in terms of emotional impact.

In the UK, PMDD may be considered a **disability under the Equality Act 2010**, as it significantly impacts an individual's ability to perform day-to-day activities. This classification is important when considering patient support and workplace accommodations.

Core characteristics:

Timing:



Symptoms typically begin during the luteal phase (1-2 weeks before menstruation) and subside almost immediately after the period starts, or within a few days after.

Severity:



PMDD symptoms are intense enough to disrupt daily life, impacting work, relationships, and overall wellbeing.

Psychological symptoms:



The emotional symptoms, especially mood swings, anxiety, and depression, are key indicators that differentiate PMDD from PMS.

COMMON SYMPTOMS OF PMDD


While symptoms vary from person to person, PMDD usually presents with a combination of emotional, cognitive, and physical symptoms.

Emotional symptoms:

- Severe mood swings
(e.g. Sudden sadness or tearfulness)
- Irritability or anger, often leading to conflicts
- Anxiety or tension, or feeling “on edge”
- Depressive thoughts or feelings of hopelessness
- Difficulty concentrating
- Fatigue or lack of energy
- Overwhelm or a sense of being out of control

Physical symptoms:

- Bloating or weight gain
- Breast tenderness
- Headaches
- Joint or muscle pain
- Sleep disturbances
(e.g. Insomnia or excessive sleep)
- Appetite changes
(e.g. Overeating or food cravings)



NEURODIVERSITY AND PMDD

There is increasing evidence that individuals who are neurodivergent, such as those with autism spectrum disorder (ASD), ADHD, or other neurodevelopmental conditions, may experience PMDD more intensely.

Neurodivergent individuals often have heightened sensitivities to sensory stimuli and emotional dysregulation, which can be exacerbated during the luteal phase of the menstrual cycle. This means that the symptoms of PMDD may not only feel more severe but can also lead to a more difficult time coping with day-to-day stressors.

Key considerations:

Sensory sensitivities:

Individuals with neurodiversity may experience amplified sensitivity to light, sound, and touch during PMDD, which can increase emotional distress.



Emotional regulation:

Those with conditions like ADHD or autism may find it more challenging to regulate emotions during the luteal phase, which can exacerbate irritability, anxiety, or mood swings.



Understanding this connection is crucial in ensuring that patients receive appropriate care and tailored support.

MAKING PATIENTS FEEL HEARD AND UNDERSTOOD

It's important to remember that your patients may downplay their symptoms due to the anxiety and self-doubt that PMDD can trigger.

They may feel embarrassed or worry that they won't be taken seriously and some patients may have had previous negative experiences with healthcare providers, so it's essential to create a supportive and non-judgemental space where they feel comfortable discussing their experience.

Ask open-ended questions:



Encourage your patients to describe how their symptoms affect their daily life, both at home and at work. This can help you gauge the true impact, even if they initially hesitate to fully express it.

Acknowledge variability:

Every patient's experience with PMDD is different. Some may have more intense mood swings, while others may experience severe physical symptoms.

Letting them know that you understand the diversity of symptoms will make them feel validated.

Reassure them they're not alone:



PMDD can make patients feel isolated or like their emotions are "too much". Reassure them that PMDD is a real, recognised condition and that they're not alone in their struggle.

DIFFERENTIATING PMDD FROM OTHER CONDITIONS

PMDD is sometimes mistaken for other mood disorders, such as bipolar disorder, depression, or anxiety.

It's crucial to differentiate PMDD by focusing on the cyclical nature of the symptoms, which follow the menstrual cycle.

PMDD vs. Bipolar disorder:

Unlike bipolar disorder, which involves sustained mood episodes that can last for weeks or months, PMDD symptoms occur predictably during the luteal phase of the menstrual cycle and resolve shortly after menstruation begins.

Patients with PMDD will have symptom-free periods between cycles.

PMDD vs. Depression/anxiety:

While PMDD can cause severe depressive or anxious feelings, these symptoms are cyclical and tied to the luteal phase. Patients with major depressive disorder or generalised anxiety disorder experience symptoms that are not linked to the menstrual cycle.

The importance of symptom tracking:

To accurately diagnose PMDD, it's essential that patients track their symptoms over at least two menstrual cycles.

Many patients may not realise the connection between their symptoms and their menstrual cycle, so encouraging them to keep a detailed symptom diary can be incredibly valuable.

Ask for daily symptom logs:



Encourage patients to record their emotional, cognitive, and physical symptoms every day for two cycles. This can help reveal patterns tied to the luteal phase.

Use symptom tracking apps:



Suggest apps specifically designed for PMDD tracking, which can help patients clearly document the timing and intensity of their symptoms. Please be aware that not all apps cater to irregular cycles.

DIAGNOSTIC CRITERIA FOR PMDD (DSM-5):

According to the DSM-5, to diagnose PMDD, a patient must experience at least five of the following symptoms during most menstrual cycles in the luteal phase, with at least one being from the first four:

Emotional/Cognitive:

Affective lability (e.g. Mood swings, sudden sadness)	Irritability or anger	Depressed mood or feelings of hopelessness	Anxiety or tension
---	------------------------------	---	---------------------------

Physical/Behavioural:

Decreased interest in usual activities	Difficulty concentrating	Lethargy or fatigue	Changes in appetite, overeating, or food cravings
Hypersomnia or insomnia	A sense of being overwhelmed or out of control	Physical symptoms such as breast tenderness, bloating, or joint pain	

Symptoms must be severe enough to interfere with daily life and not solely attributable to another mental health disorder. In some cases, you may need to refer the patient to a psychologist or psychiatrist for an official diagnosis or further mental health support, as PMDD often requires multidisciplinary approach to treatment.

In the UK, psychologists or psychiatrists can help with a more formal mental health evaluation, and psychiatrists can prescribe medications if needed.

Offering compassionate care:

Understanding the emotional toll PMDD can take on your patients is essential for providing compassionate care.

Let them know that you take their symptoms seriously and that you'll work with them to find an effective management plan.

Discuss treatment options:



Options can range from lifestyle modifications and dietary changes to hormonal treatments, antidepressants, or Cognitive Behavioural Therapy (CBT). Collaborate with your patient to find the best treatment plan for their individual needs.

Encourage self-advocacy:



Empower patients to speak up about their needs and seek support from family, friends, or mental health professionals.

Some patients may also be entitled to workplace accommodations.

Regular follow-ups:



Schedule regular check-ins with patients to monitor their progress and make adjustments to their treatment plan as needed.

As a GP, your attentiveness and empathy can make a profound difference in helping patients with PMDD. Recognising the cyclical nature of the disorder and offering a supportive, non-judgemental environment will ensure patients feel heard and understood.

Given that PMDD can be recognised as a disability under the Equality Act 2010, ensuring patients have the right support at work and in their personal lives is essential.

With your help, they can access the right treatment and regain control over their health and wellbeing.

Disclaimer: Everyone's menstrual cycle is unique, and not all cycles last exactly 28 days.

The information provided in this resources is meant to offer general guidance, but it's important to remember that your cycle length and phases may vary. Some cycles may be shorter or longer, and the intensity of symptoms and energy levels can differ from person to person. Always listen to your body and consult with a healthcare professional to tailor any advice to your individual needs.

THE PMDD PROJECT